

PHA 5127
Answers Case Study 2
Fall 2007

1. Patient A.M. is given a 60 mg dose of gentamicin. The volume of distribution for this patient is 10 L and the concentration after 8 hours is 1.4 mg/L. Calculate the k_e . What is the half-life?

Answer: $C_0 = \text{Dose}/(\text{Volume of distribution}) = 60\text{mg}/10\text{L} = 6\text{mg/L}$
 $C = C_0 * e^{-k_e * t}$ $(\ln C - \ln C_0) = -k_e * t \rightarrow -(\ln C - \ln C_0)/t = k_e \rightarrow k_e = -(\ln 1.4\text{mg/L} - \ln 6\text{mg/L})/8 \text{ hours}$
 $\rightarrow k_e = 0.182 \text{ hours}^{-1}$

$t_{1/2} = 0.693/k_e \rightarrow t_{1/2} = 0.693/0.182 \text{ hours}^{-1} = 3.81 \text{ hours}$

2. Assuming a one compartment body model and a 1st order process, please graph the following on semilog paper and predict the concentration after 6 hours.

Time (hours)	Concentration (mg/L)
1	80
3	42
5	22

Answer = ~16 mg/L

3. Using 110 mg/L as the starting concentration and a k_e of 0.318 hour^{-1} calculate the concentration after 6 hours.

Answer: $C = C_0 * e^{-k_e * t}$ $C = 110 \text{ mg/L} * e^{-(0.318\text{hour}^{-1} * 6\text{hour})} = 16.3 \text{ mg/L}$

4. True or False

a. In a one-compartment body model it is assumed that a drug distributes to all areas of the body instantaneously. True.

b. Pharmacodynamics is the study of the time course of a drug's absorption, distribution, metabolism, and elimination. False.

Pharmacodynamics refers to the relationship between concentration at the site of action and the resulting effect.

c. The k_e of a drug is 0.00333 min^{-1} . After 2 hours 67% of the drug is remaining in the body. True.

Answer: $e^{-k_e \cdot t} = \text{fraction remaining}$ $e^{(-0.00333 \text{ min}^{-1} \cdot 60 \text{ min/hour} \cdot 2 \text{ hours})} = 0.67$

